

AMENDED SPECIFICATION

54

RECEIVED

JUL 12 2001

TECH CENTER 1600/2900



Example 4

Construction of RADs to prevent *Streptococcus equi* infection and disease.

- 10 *Streptococcus equi* causes a very severe disease of racehorses and other equines called
strangles (Nara, P. et al. (1983) Am. J. Vet. Res. 44:529-534). Even worse is the immune
complex disease *Purpura hemorrhagica*, a consequence of earlier in life *S. equi* infection
(Galan, J. and J. Timoney (1985) J. Immunology 135:3134-3137). *S. equi*, like group A
streptococci, produce an M protein on the surface which is antiphagocytic and thus a major
15 virulence determinant enhancing the success of *S. equi* infection. Antibodies to the *S. equi* M
protein (SeM) are opsonic and facilitate the successful phagocytosis and killing of *S. equi*
(Galan, J. and J. Timoney (1985) Infect. Immun. 47:623-628). Thus an antibody response to
SeM is likely to be highly protective in preventing *S. equi* infection of horses. The RAV
pMEG-573 encoding the *S. equi* SeM protein is depicted in Figure 18. This RAV was
20 obtained by cloning the PCR fragment flanked by primers SeM444-474
GCGAACTCTGAGGTTAGTCGTACGGCGACTC (SEQ ID NO:1) and SeM1265-1233
TTGATCAATTTCTGCTAATTTTGTAGCCATTTC (SEQ ID NO:2), containing the central
portion of the SeM coding region from the SeM clone pSEM06, into the *NcoI* and *BamHI*
sites of pMEG-546. pMEG-573 is only dependent on the presence of the
25 $\Delta ilvG3::TTaraCP_{BAD}lacITT$ deletion/insertion mutation (Figure 2) in the chromosome to
repress the runaway phenotype and SeM expression. The vaccine strains for SeM also
contain either the $\Delta phoP1918$ or $\Delta phoP24$ attenuating deletion mutation (mutations 8 and 9,
Figure 2). A comparison of the level of SeM expression by different attenuated *Salmonella*
vaccine strains, in which SeM